


SYPHILIS TESTING ALGORITHM

Patient History & Physical Exam*

For Pregnant Patients: Screen for syphilis three times every pregnancy (first encounter, 28 weeks, and delivery)



Treating syphilis in pregnant people:

- Penicillin G benzathine (i.e., Bicillin-LA) is the only proven effective treatment for syphilis during pregnancy.
- If the patient has a true penicillin allergy as confirmed by questioning and possible evaluation by an allergy specialist, arrange for immediate desensitization and treatment.

CIA, chemoluminescence immunoassay; EIA, enzyme immunoassay; POC, point of care; RPR, rapid plasma reagin; TP-PA, Treponema pallidum particle agglutination assay; VDRL, Venereal Disease Research Laboratory test.
 * If a person has signs and/or symptoms consistent with primary or secondary syphilis, consider immediate presumptive treatment. People exposed through sexual contact to a person who has syphilis should also be considered for immediate presumptive treatment based on CDC STI Treatment Guidelines (<https://www.cdc.gov/std/treatment-guidelines/syphilis.htm>)
 † If patient has had sexual activity with someone who might have syphilis in the last 3 weeks, consider repeat syphilis testing in 2-4 weeks.
 ‡ Compare to previous RPR as available and treat per guidelines.

Copyright October 2024 American College of Obstetricians and Gynecologists
 This information is designed as an educational resource to aid clinicians in providing care, and use of this information is voluntary. This information should not be considered as inclusive of all proper treatments or methods of care or as a statement of the standard of care. This information does not represent ACOG clinical guidance. It is not intended to substitute for the independent professional judgment of the treating clinician. For ACOG's complete disclaimer, visit www.acog.org/iz-disclaimer.

